

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/340122235>

Virtual Screening of Inhibitors Against Spike Glycoprotein of SARS-CoV-2: A Drug Repurposing Approach

Preprint · March 2020

DOI: 10.20944/preprints202003.0042.v2

CITATIONS

0

READS

16

3 authors:



Kanishka Senathilake
University of Colombo

9 PUBLICATIONS 25 CITATIONS

[SEE PROFILE](#)



Sameera Ranganath Samarakoon
University of Colombo

67 PUBLICATIONS 342 CITATIONS

[SEE PROFILE](#)



Kamani Tennekoon
University of Colombo

122 PUBLICATIONS 714 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Anti-cancer effects of *Mangifera zeylanica* [View project](#)



Anti-cancer natural product development [View project](#)

Virtual screening of inhibitors against spike glycoprotein of SARS-CoV-2: a drug repurposing approach

Kanishka S Senathilake¹, Sameera R Samarakoon¹, Kamani H Tennekoon¹

Institute of Biochemistry, Molecular Biology and Biotechnology, University of Colombo, Colombo 03. Sri Lanka.

Author of correspondence - Sameera R Samarakoon email - sam@ibmbb.cmb.ac.lk

Abstract: The novel coronavirus (SARS-CoV-2) is a human pathogen recently emerged in China, causing a global pandemic of severe respiratory illness (COVID19). SARS-CoV-2 makes entry in to human cells through its spike (S) protein that binds to cell surface receptors. Widespread of SARS-CoV-2 has been attributed to high affinity of S protein to its receptor. A homology model of the receptor binding domain of SARS-CoV-2 S protein (RBD) was built. RBD- receptor docking and published molecular dynamics data were used to map the key RBD-receptor interaction hotspot (RBDhp) on the RBD. Primary virtual screening was carried out against RBDhp using more than 3300 compounds approved by U.S Food and Drug Administration (FDA) and other authorities for human use. Compounds that bind to hpRBD with a binding energy $\leq - 6.5$ kcal/mol were subjected to secondary screening using a recently published cryo EM (2.9 Å) structure of RBD. A cardiac glycoside (digitoxin), two anthracyclines (zorubicin and aclarubicin), a tetracycline derivative (rolitetracycline), a cephalosporin (cefoperazone) and a food dye (E-155) were predicted to be most potent inhibitors of RBD – receptor interaction. An anti-asthmatic drug (zafirlukast) and several other drugs (itraconazole, fentanyl, troglitazone, gliquidone, Idarubicin, Oxacillin) were found to be high affinity binders that may have a potential to inhibit RBD – receptor interaction.

Keywords: Flue; Virus; COVID19; SARS CoV-2; Spike Protein; Virtual Screening; digitoxin, anthracycline, rolitetracycline, E-155

1. Introduction

The outbreak of coronavirus disease (COVID-19) which was first reported from Wuhan, China has been spreading around the world [1]. World Health organization has declared the SARS-CoV-2 as a Public Health Emergency of International Concern (PHEIC) on the 30th of January 2020 by considering its rapid transmission from human to human and spread into many of the continents [2-3]. It has now been recognized by WHO as a global pandemic making it the first pandemic caused by a coronavirus [4]. Currently the number of confirmed infections around the world stands at 167 511 while a total of 6606 lives have been claimed since its initial outbreak in December 2019 [4].

Currently there are more than 350 genome sequences of SARS-CoV-2 that have been shared on the online platform Global Initiative on Sharing All Influenza Data (GISAID). The 30,000 base-pair genome of SARS-CoV 2 shows that the virus is adequately divergent from SARS-CoV; a human-infecting beta-coronavirus that caused an epidemic in 2003 [5]. Further, based on the genome variations located in open reading frame 8 (for which the function is yet unknown), Tang and others have identified two viral lineages which are named as L and S which are likely to be evolved in early stages of outbreak in china [6]. Like other viruses, SARS-CoV-2 evolves over time introducing random mutations. While the mutation rate for SARS-CoV-2 has not been calculated accurately it has been suggested to have moderate to high mutation rate when compared to other members of single-stranded RNA viruses [7].

SARS-CoV-2 binds to the human angiotensin converting enzyme 2 (ACE-2) receptor through densely glycosylated spike (S) protein as the initiation step of the entry mechanism to human cells [8-9]. The S protein of SARS-CoV-2 binds ACE-2 through its receptor binding domains (RBD) and the RBD – up conformation of the S protein is a prerequisite for the formation of RBD-ACE-2 complex [9]. Divergence of SARS-CoV-2 RBD amino acid sequence from other corona viruses and RBD-ACE-2 binding experiments has given implications on relatively high affinity viral binding to host cell receptors [9].

High affinity RBD-ACE-2 binding has been suggested among other factors for wide spread of SARS-CoV-2. Although highly important drug target, unavailability of a high-resolution crystal structure and solvent accessible binding surface has made it a tedious target for virtual screening and drug design[10]. Cryo EM structure of SARS-CoV-2 S protein (pre fusion down conformation) has been published at 3.5 Å resolution [9]. Availability of several high

resolution X-ray crystal structures of SARS CoV S protein (UP conformation) in complexed with ACE-2, structural rigidity and high degree of sequence similarity of RBDs warrants us to generate reasonably accurate homology models for drug screening. In the current study more than 3000 compounds approved by various regulatory authorities including U.S. Food and Drug Administration (FDA) were screened against an optimized homology model of the SARS-CoV-2 S protein RBD. After publishing our initial work in the first version of the preprint [11] a 2.9 Å cryo EM structure (PDB 6m17) of RBD-ACE-2 complex have been published [12]. High resolution of 6m17 and low temperature factor of the RBD-ACE-2 interface warranted carrying out a second round of screening using the new cryo EM structure.

2. Results and discussion

2.1. Molecular modeling

Spike protein was selected for virtual screening instead of selecting ACE -2 receptor since compounds that block ACE -2 receptor is known to have modulatory effect on blood pressure and several other cardiovascular system related side effects [13]. Cryo EM map of SARS-CoV-2 S protein has been published in receptor unbound pre fusion conformation at a resolution of 3.5 Å [9]. The Cryo EM structure reflects the high degree of structural homology between SARS-CoV-2 RBD and SARS- CoV RBD having a RMSD value of 3.0 Å [9]. Low resolution and the conformational considerations of Cryo EM structure published by Wrapp et al, makes it an unreliable structure for virtual screening [9]. Therefore, a homology model was generated using high-resolution crystal structure of the SARS -CoV (PDB ID:2AJF) in ACE-2 bound up confirmation. Energy minimized homology model used for the screening showed a C α RMSD of 1.47 Å between the RBD domains of SARS-CoV-2 and the SARS-CoV.

2.2. Identification of Ligand Binding Site

Analysis of interaction surface of RBD-ACE2 in chimera showed strong H bonding at TYR 505 which is well exposed indicating an initial contact point with ACE-2. Further polar interactions were observed at THR 500, ASN 501, GLY 502, TYR 449, GLN 493 and GLN 498. This approach gives interactions in consistence with pervious protein - protein docking data [14] and data arising from MD experiments [15]. Further surface analysis of the structure revealed Arg 439 > ASN 439 mutation on one side of the structure that is protruding

(SpkP) from the rest of the S protein (Figure 1). This mutation seems to significantly weaken the hydrogen bond formed by residue number 439. Therefore, the key interactions were restricted to one side of SpkP. (Figure 1). For THR 500, ASN 501, GLN 498, observations were in consistency with the RBD-ACE-2 interactions observed in the most recent publication of cryo EM structure 6m17 [12]. A comparison of binding residues is given in figure 1.

We took the advantage of ARG439> ASN 439 mutation in SARS-CoV-2 to define a potent inhibitor binding site on the side opposite to ASN 439. The ligand binding site spanned from GLN 498 through GLN 493 and Tyr 505 and ending in a hydrophobic pocket consisting LEU 555, ILE 418 and hydrophobic regions of LYS 417, TYR 421 and TYR 453. (Figure 1).




Figure 1- Ligand binding site (hpRBD, white color dotted circle) on the SARS-CoV-2 receptor binding domain (RBD) used for virtual screening. a. Homology model of RBD (RBD-HM, template- 2AJF). b. Cryo EM structure of RBD at 2.9 Å resolution (6m17). Red color residues are the previously identified key binding residues [11,15]. Magenta color residues are the residues that form polar RBD-ACE-2 interactions observable in 6m17. Red dotted circle shows the open hydrophobic pocket of BRD-HM which is closed in 6m17.

2.3. Virtual Screening

Protein ligand interaction analysis was carried out for hits (binding energy \leq -6.5 kcal/mol) resulting from secondary screening. According to our previous experience with protein-

protein interaction inhibitors, a two hit hypothesis (targeting at least two key residues that form polar proteins – protein interactions) was employed to select potent inhibitors of S protein and ACE-2. This approach when coupled with high energy binding, provides a better approach for inhibiting relatively strong protein-protein interactions that span through large surface area. Detailed list of interactions formed by the hits identified in the current study are given in table 1.

Digitoxin, (ZINC accession - 8101077) a FDA approved cardiac glycoside formed hydrogen bonds with four residues critical in forming RBD-ACE-2 interaction (table 1, figure 3a). This lowest energy pose had a binding energy value of **-7.6 kcal/mol** indicating its potential as a RBD-ACE-2 interaction inhibitor. Digitoxin is used in the treatment and management of cardiac congestion, arrhythmias and heart failure (used in place of digoxin) [16]. Use of digitoxin is decreasing owing to the availability of better replacement drugs and some toxic properties (anorexia, nausea and vomiting) [16-17].

The first and second binding poses and fifth binding pose of zorubicin (ZINC accession - 03831623) exhibited interesting interactions (table 1, figure 3 b-d). In particular, the pose one was observed to form hydrogen bonds with Gln 498 and Thr 500 that have been experimentally elucidated to form most important strong polar interactions with ACE-2 [12]. Thus the compound picks two most critical residues through hydrogen bonds while forming hydrophobic interactions with Gln493 and Tyr 505. Zorubicin is classified in the zinc library as a world drug (not approved by FDA) and it is approved in several European countries including France for the treatment of acute leukemia [18].

Aclarubicin (ZINC accession - 08101054) is an anthracycline drug (world drug not approved by FDA) that is similar to zorubicin. It is used with cytarabine in Japan, China and India for the treatment of different acute myeloid leukemia conditions [19]. Hydrogen bonds were observed at Gln 498 and Try 453, Gln493 and Tyr 449 in the lowest energy binding pose of aclarubicin. Thus we provide strong evidence that the aclarubicin is a potent candidate for the inhibition of RBD -ACE-2 interaction. (Table 1). However anthracycline drugs such as zorubicin and aclarubicin are known to cause cardiac toxicity, inhibition of DNA replication and repair when used for a longer period of time [18,20]

Rolitetracycline (ZINC accession - 3831437) is a FDA approved broad-spectrum tetracycline antibiotic particularly used for parenteral administration in cases requiring elevated concentrations. At the therapeutic concentrations rolitetracycline does not cause toxic effects.

[21]. The first and the second lower energy binding poses of rolitetracycline interacted with at least two key residues forming hydrogen bonds (table 1, figure 3 e). In both poses, strong hydrogen bonds were observed with Tyr 453. Therefore rolitetracycline is a potent candidate for the inhibition RBD -ACE-2 interaction.

E-115 (Brown HT) is a common food dye (bis-azo dye) authorized as a food additive in the European Union (EU) [22]. Lowest energy poses showed hydrogen bonding with key residues Gln 498 and Asn 501 while the second lowest energy binding pose showed hydrogen bonding with key residues Thr 500 and Asn 501. Thus the current study provides evidence for E 155 as a potent candidate for the inhibition RBD -ACE-2 interaction. (Table 1, figure 3 f).

Cefoperazone is a FDA approved beta-lactam antibiotic used in the treatment of various bacterial infections including respiratory tract infections. While the clinical use of cefoperazone is discontinued in the US, it is commonly available in several European countries for therapeutic use. First, second and the fourth binding poses of cefoperazone showed hydrogen bonding with key residues. Lowest energy poses showed hydrogen bonding with key residues Tyr 449 and Gln 493 while the second lowest energy binding pose showed hydrogen bonding with Gln 493 and Asn 501. In the fourth binding pose, hydrogen bonds were observed with Lys 417, Tyr 453, Gln 493 and Asn 501. (Table 1, figure 3 g) indicating the potential of cefoperazone as a potent RBD -ACE-2 interaction inhibitor.

Interestingly a commonly used anti asthmatic drug (zafirlukast) was predicted to bind with high affinity (-7.4 kcal/mol). Zafirlukast forms hydrogen bond with only one key residue (applies to all poses) while in all the poses zafirlukast stands in the binding site giving steric hindrance to key residues (not shown). Therefore, considering the high affinity and its common use as an asthmatic drug, zafirlukast may be tested on COVID 19 patients without waiting for in vitro data.

Further, an azole antifungal (itraconazole), a phenylimidazole muscle relaxant (tizanidine), a 1-benzopyran antidiabetic drug (Troglitazone), an antidiabetic sulfonylurea drug (gliclazide), an anthracycline antineoplastic drug (Idarubicin) and a beta-lactam antibiotic (Oxacillin) were found to be high affinity binders that may have the potential to inhibit RBD – receptor interaction (table 1).

Table 1. **Compounds that bind (binding energy ≤ -6.5 kcal/m^{ol})** and position within the ligand binding site (RBDhp). BE = binding energy (pose number is given in brackets), HB = hydrogen bonding. HP = Hydrophobic interaction, $\Pi\Pi$ = phi-phi stacking. Lengths of hydrogen bonds are given as donor -acceptor distance (>3.5 Å bonds are shown in gray color). Common names were retrieved from without considering chiral nature.

	Zinc accession Number	Common name (Drug Bank/ PubChem)	Binding energy (kcal/m ^{ol})	Interactions with key binding site residues and bond lengths (Å)								
				Lys 417	Tyr 449	Tyr 453	Gln 493	Gln 498	Thr 500	Asn 501	Tyr 505	
Compounds that form HB with ≥ 2 key residues	8101077	Digitoxin	-7.6(0)	-	3.0	-	3.9	-	-	3.6	3.0	
	03831623	Zorubicin	-7.6(0)	-	4.0	-	2.2	-	-	-	-	
			-7.4(1)	-	3.9	-	HP	2.9	3.1	4.0	HP	
			-7.2(5)	-	3.9	2.7	2.9	3.6	-	3.7	-	
	08101054	Aclarubicin	-6.6(0)	-	2.8	3.3	3.3	2.9	-	-	-	
	03831437	Rolitetracycline	-6.6(0)	HP	-	2.9	-	-	-	-	-	-
			-6.6(1)	-	2.6	2.9	2.3	-	-	2.6	-	
	03830332	E155	-7.9(0)	-	-	-	-	3.2	-	3.9	$\Pi\Pi$ S	
			-7.6(1)	-	-	-	-	-	3.8	3.1	$\Pi\Pi$ S	
	03830429	Cefoperazone	-7.0(0)	-	2.8	-	3.1	-	-	-	$\Pi\Pi$ S	
			-6.8(1)	-	-	3.8	3.0	-	-	3.3	$\Pi\Pi$ S	
			-6.6(4)	1.9	-	2.8	2.1	-	-	2.4	2.9	
03830973	Itrazol	-6.6(3)	-	-	3.0	-	-	3.0	-	3.0		
00896717	Zafirlukast	-7.4(0)	-	-	-	2.8	-	-	-	-		
03830826	Fazadinium	-7.4(0)	-	-	-	-	HP	-	3.2	$\Pi\Pi$ S		
		-7.4(1)	-	-	-	-	HP	-	3.2	$\Pi\Pi$ S		
01482077	Gliquidone	-7.0(4)	-	-	-	2.6	-	-	-	-		
00968277	Troglitazone	-6.8(2)	-	-	3.3	-	-	-	-	-		
00968279	Troglitazone	-6.7(4)	-	-	-	2.9	-	-	-	-		
11678097	Idarubicin	-6.7(0)	-	-	-	-	-	3.2	-	-		
01530747	Oxacillin	-6.6(0)	-	-	-	-	2.3	-	-	-		







Figure 2. Structures and chirality of inhibitors that form at least two hydrogen bonds with key target residues of RBD. a. Digitoxin (8101077), b. Zorubicin (03831623), c. Aclarubicin (08101054) d. E155 (03830332), e. Rolitetracycline (03831437) f. Cefoperazone (03830429)




a




b




c




d




e



f



g



h




Figure 3. Binding poses of compounds listed in table 1, a. digitoxin (0), b-d. zorubicin (0,1,5), e-f. Rolitetracycline (0.1), g. Aclarubicin (0), h-i. E115 (0.1) j-l. cefoperazone.

3. Materials and method

3.1. Compound library, software and hardware.

The compound library (3350 compounds) was originally prepared for FightAIDS@Home project by Stefano Forli at Scripps Research Institute (TSRI, La Jolla, CA) [23]. The library containing geometry optimized 3D structures of all the tautomers and enantiomers of approved compounds (FDA and other world authorities) was downloaded from Zinc database (<http://zinc.docking.org/pdbqt/> accessed in December 2017) and available on the local server for screening.

Swiss Model [24] was used to generate homology models. For virtual screening, autodock vina program [25] in the academic distribution of PyRx [26] was run in the local server computer (Intel Xeon CPU E5-2420 v2, 2.2 GHz, 12 core processor, 32 GB RAM, Ubuntu OS 16.04 LTS). For protein structure manipulation, comparison and binding pose analysis,

Chimera v 1.14 [15] was run on a personal computer (Intel® Core™ i7-3540M processor ,4 GB ram, windows 10). Protein Ligand Interaction Profiler (PLIP) web server [27] was used for detailed analysis of protein-ligand complexes.

3.2. Molecular modeling

Homology models of the RBD was constructed using the crystal structure (2.9 Å) of RBD of SARS CoV (RBD domain complexed with ACE2 (PDB: 2AJF_E) as the template (76.47% sequence identity.). As previous studies have shown to preserve key interactions between RBD and ACE-2 of SARS CoV and SARS-CoV-2, RBDs of SARS-CoV-2 (homology model) and SARS CoV (2AJF_E) were superimposed on RBD of SARS CoV using match maker function of chimera. Resulted RBD – ACE-2 complex was then energy minimized using chimera (400 steps of steepest descents, 50 steps of conjugated descents) while keeping the backbone of the ACE-2 fixed [28,29] using general Amber force field [30] to obtain the final homology model of the RBD (RBD-HM). Instead of using conformational ensembles derived from complete molecular dynamic (MD) simulations or structures derived from simulated annealing [31], the single model generated using the afore mentioned approach was selected for virtual screening as it provides adequate evidence for a reliable model for virtual screening (see section 2).

3.3. Primary Screening

Autodock Vina Grid Box was set to cover all the residues of RBD-HM that form key interactions with ACE-2 (center X= 166.39, Y = 170, 405, Z = 135.72, dimensions X = 28.91 Y = 28.59, Z = 28.55). Finally, all the ligands were docked using the Lamarckian genetic algorithm as scoring function. TYR 449, GLN 493, GLN 498, THR 500, TYR 501 and TYR 505 were made flexible and the other parameters were default values.

3.4. Secondary Screening

After carrying out the primary screening, ligands having binding energy values less than -6.5 kcal/mol were used for secondary screening against 6m17. For the secondary screening the RBD of 6m17 chain E was retained and other polypeptide chains and water molecules were removed. Screening was carried out using the same method as described in section 2.2. Binding poses having binding energy vale \leq -6.5 kcal/mol were characterized using view dock function in chimera and PLIP web server to identify the profile of protein -ligand

interactions. Compounds that fit in to the binding site and form polar interactions with at least two critical residues were considered as potent inhibitors of RBD-ACE-2 interaction.

4. Conclusion

In conclusion, a cardiac glycoside (digitoxin), Four antibiotics; two anthracyclines (zorubicin and aclarubicin), a tetracycline derivative (rolitetracycline) and a cephalosporin (cefoperazone), a food dye (E-155) were predicted to be potent inhibitors of RBD – receptor interaction based on the selection criteria used. A commonly used anti-asthmatic drug (zafirlukast) and several other drugs (itrazole, fazadinium, troglitazone, gliquidone, idarubicin, oxacillin) were found to be high affinity binders that may have potential to inhibit RBD – receptor interaction. Results of present study suggest the potential of these compounds as prophylactic medication or use in preventive countermeasures.

5. Compliance and ethics

The authors declare that they have no conflict of interest.

6. Acknowledgement

Mr. Kanchana Senanayake. Assistant network manager of the Institute of Biochemistry, Molecular Biology and Biotechnology for providing computing facility

7. References

1. Chan JF, Yuan S, Kok KH, To KK, Chu H, Yang J, Xing F, Liu J, Yip CC, Poon RW, Tsoi HW, Lo SK, Chan KH, Poon VK, Chan WM, Ip JD, Cai JP, Cheng VC, Chen H, Hui CK, Yuen KY. (2020) A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*. 395:514-523
2. Sun K, Chen J, Viboud C. Early epidemiological analysis of the coronavirus disease 2019 outbreak based on crowdsourced data: a populationlevel observational study. *Lancet Digital Health* 2020 Published Online February 20, 2020 [https://doi.org/10.1016/S2589-7500\(20\)30026-1](https://doi.org/10.1016/S2589-7500(20)30026-1).
3. World Health Organization (a), Coronavirus disease 2019 (COVID-19) Situation Report – 38. 27.02.2020. Accessed online on 28th February 2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200227-sitrep-38-covid-19.pdf?sfvrsn=9f98940c_2
4. World Health Organization (b), Coronavirus disease 2019 (COVID-19) Situation Report – 56.16.03. 2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200316-sitrep-56-covid-19.pdf?sfvrsn=9fda7db2_2

5. Lu R, Zhao X, Li J, Niu P, Yang B, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* 2020; 395: 565–74.
6. Tang X, Wu C, Li X, Song Y, Yao X, Wu X, Duan Y, Zhang H, Wang Yi, Qian Z, Cui J, Lu J, On the origin and continuing evolution of SARS-CoV-2, *National Science Review*, nwa036, <https://doi.org/10.1093/nsr/nwaa036>
7. Amodio E, Vitale F, Cimino L, Casuccio A, Tramuto F, Outbreak of Novel Coronavirus (SARS-Cov-2): First Evidences From International Scientific Literature and Pending Questions *Healthcare* 2020, 8, 51; doi:10.3390/healthcare8010051
8. Liu C, Zhou Q, Li Y, Garner L.V, Watkins S.P, Carter, L.J, Smoot J, Gregg A C, Daniels A D, Jervey S, Albaiu D. Research and Development on Therapeutic Agents and Vaccines for COVID-19 and Related Human Coronavirus Diseases. *ACS Central Science*. 2020;. doi:10.1021/acscentsci.0c00272
9. Wrapp D, Wang N, Corbett K S., Goldsmith J A., Hsieh C, Abiona O, Graham B S, McLellan J S. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation *Science* 19 Feb 2020:DOI: 10.1126/science.abb2507
10. Gruber C, Steinkellner G: Coronavirus COVID-19 (formerly known as Wuhan coronavirus and SARS-CoV-2) - what we can find out on a structural bioinformatics level. 2020.
11. Senathilake, K.; Samarakoon, S.; Tennekoon, K. Virtual Screening of Inhibitors Against Spike Glycoprotein of 2019 Novel Corona Virus: A Drug Repurposing Approach. Preprints 2020, 2020030042 (doi: 10.20944/preprints202003.0042.v1)
12. Wan, Y., Shang, J., Graham, R., Baric, R. S., & Li, F. (2020). Receptor recognition by novel coronavirus from Wuhan: An analysis based on decade-long structural studies of SARS. *Journal of Virology*. doi:10.1128/jvi.00127-20
13. Mohammed A.R ,Chamsi-Pasha MA, Shao Z, Tang WH. Angiotensin-converting enzyme 2 as a therapeutic target for heart failure. *Curr Heart Fail Rep*. 2014;11(1):58–63. doi:10.1007/s11897-013-0178-0.
14. Xu X, Chen v, Wang J, Feng J, Zhou H, Hao L X, W Z P Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission, 2020. *Sci China Life Sci*. 2020 Jan 21. doi: 10.1007/s11427-020-1637-5.
15. Peng, Cheng; Zhu, Zhengdan; Shi, Yulong; Wang, Xiaoyu; Mu, Kaijie; Yang, Yanqing; et al. (2020): Exploring the Binding Mechanism and Accessible Angle of SARS-CoV-2 Spike and ACE2 by Molecular Dynamics Simulation and Free Energy Calculation. ChemRxiv. Preprint. <https://doi.org/10.26434/chemrxiv.11877492.v1>

16. Lipman, A.G. (1993), Martindale: 'Martindale — the Extra Pharmacopoeia' (30th ed), edited by J. E. F. Reynolds. International Journal of Pharmacy Practice, 2: 124-124. doi:10.1111/j.2042-7174.1993.tb00740.x
17. Johan H. Digitoxin has Specific Properties for Potential use to Treat Cancer and Inflammatory Diseases. Res & Rev Health Care Open Acc J 2(3)- 2018. RRHOAJ.MS.ID.000137.
18. Hiddemann W, Büchner T, Plunkett W, Keating M, Wörmann B, Andreeff M. Acute Leukemias: Pharmacokinetics and Management of Relapsed and Refractory Disease. Springer Science & Business Media publishing (2012), 670 pages
19. Cresteil, T. Aclarubicin. Reference Module in Biomedical Sciences. 2017; doi:10.1016/b978-0-12-801238-3.97571-8
20. Edwardson DW, Narendrula R, Chewchuk S, Mispel-Beyer K, Mapletoft JP, Parissenti AM. Role of Drug Metabolism in the Cytotoxicity and Clinical Efficacy of Anthracyclines. Curr Drug Metab. 2015;16(6):412–426. doi:10.2174/1389200216888150915112039
21. Griffin MO, Fricovsky E, Ceballos G, Villarreal F: Tetracyclines: a pleiotropic family of compounds with promising therapeutic properties. Review of the literature. Am J Physiol Cell Physiol. 2010;299(3):C539-48. doi: 10.1152/ajpcell.00047.2010.
22. EFSA Journal 2014;12(5):3719. 26 pp. doi:10.2903/j.efsa.2014.3719. Available online: www.efsa.europa.eu/efsajournal
23. Perryman AL , Forli S Dallakyan S, Morris GM, Olson AJ, FightAIDS@Home newsletter (2009) 8: <http://fightaidsathome.scripps.edu/images/FAAHvol8.pdf>
24. Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., Heer, F.T., de Beer, T.A.P., Rempfer, C., Bordoli, L., Lepore, R., Schwede, T. SWISS-MODEL: homology modelling of protein structures and complexes. Nucleic Acids Res. 46, W296-W303 (2018).
25. Trott O, Olson A J, AutoDockVina: improving the speed and accuracy of docking with a new scoring function, efficient optimization and multithreading, Journal of Computational Chemistry 31 (2010) 455-461
26. Dallakyan S, Olson AJ. Small-molecule library screening by docking with PyRx. Methods Mol Biol. 2015;1263:243-50
27. Salentin S, Schreiber S, Haupt VJ, Adasme MF, Schroeder M (2015) PLIP: Fully automated protein-ligand interaction profiler. Nucleic Acids Res 43(W1):W443–W447

28. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. J. UCSF Chimera--a visualization system for exploratory research and analysis. *Comput Chem.* 2004 Oct;25(13):1605-12.
29. Yang Z, Lasker K, Schneidman-Duhovny D, Webb B, Huang CC, Pettersen EF, Goddard TD, Meng EC, Sali A, Ferrin TE. UCSF Chimera, MODELLER, and IMP: An integrated modeling system. *J Struct Biol.* 2012 Sep;179(3):269-78
30. James A. Maier, Carmenza Martinez, Koushik Kasavajhala, Lauren Wickstrom, Kevin E. Hauser, and Carlos Simmerling. ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *Journal of Chemical Theory and Computation* 2015 11 (8), 3696-3713 DOI: 10.1021/acs.jctc.5b00255.
31. Sali A & Blundell T.L. Comparative protein modelling by satisfaction of spatial restraints. *J. Mol. Biol.* 234, 779-815, 1993.

